

**M. SC. PHYSICS EXAMINATION, 2023**

( 2nd Year, 2nd Semester )

( 3rd Year, 2nd Semester )

**PLASMA PHYSICS**

**PAPER – 301**

Time : 2 hours

Full Marks : 40

*The figures in the margin indicate full marks.*

*Candidates are instructed to give their answers in their own words as far as practicable.*

Answer any **four** questions.  $4 \times 10 = 40$

1. Plasma is a collection of charge particles (mainly electrons and ions) in which collective effects dominate due to long range Coulomb forces. One example of plasma collective behaviour is 'Debye shielding'. Consider a plasma is in equilibrium, and we introduce a test charge  $Q_T$  inside the plasma.
  - a) Explain physically the effects of the test charge  $Q_T$  on the equilibrium plasma particles.
  - b) Prove that the potential due to the test charge  $Q_T$  obeys the formula  $\phi(r) = \frac{Q_T}{4\pi\epsilon_0 r} e^{-r/\lambda_D}$ , where  $\lambda_D = \sqrt{\epsilon_0 k_B T_e / n_0 e^2}$  is known as Debye length and all other symbols have their usual meanings.
  - c) Roughly sketch a  $\phi(r)$  vs.  $r$ , curve clearly showing whether the potential due to the test charge falls off much faster in plasma than in vacuum or not.

2+6+2

[ Turn over

[ 2 ]

2. In the context of motion of charged particles in uniform electric and magnetic fields ( $\vec{E} \perp \vec{B}$ ), we have learnt that both the ions and electrons experience  $\vec{E} \times \vec{B}$  drift with same drift velocity  $\vec{v}_d = \frac{\vec{E} \times \vec{B}}{B^2}$ . Consider a plasma is immersed in an external uniform magnetic field which points along the  $z$  direction in a Cartesian coordinate system, i.e,  $\vec{B} = B_0 \vec{k}$ . Moreover, we now take the electric field to be non-uniform and it is along the  $x$  direction, i.e.  $\vec{E} = E_0 \cos(\alpha x) \hat{i}$ , where  $\alpha^{-1}$  is the scale length of variation of non-uniform electric field.

- a) Prove that, in the presence of such a non-uniform electric field, the above expression of the drift velocity will be modified to give  $\vec{v}_d = \frac{\vec{E} \times \vec{B}}{B^2} \left( 1 - \frac{1}{4} \alpha^2 r^2 \right)$ , where the second term in the first bracket is called the finite Larmor radius effect, with  $r$  denotes the Larmor radius.
- b) Hence physically explain what will happen if somehow a density clump occurs inside the plasma due to the presence of a non-uniform electric field.
- c) Does the plasma drive any current in the presence of the non-uniform electric field and uniform magnetic field? 7+2+1

[ 5 ]

6.
  - a) What is the necessity of vacuum during the plasma formation for material processing?
  - b) How plasma diagnostics can help to optimize the material properties in a shorter time?
  - c) For high resistive material processing which type of plasma between DC and RF is preferred and why?
  - d) The following graph (see Figure 1) shows the effect of radio frequency (RF) power on the deposition rate for a sputtering process. Describe the nature of the curve with possible explanations. Why deposition rate decreased at higher RF power? 2+3+2+3
7.
  - a) What is the basic difference between plasma deposition and plasma etching? Name some processes of plasma deposition and plasma etching.
  - b) What are the input control parameters for the formation of glow discharge plasma during semiconductor material deposition? How input parameters affect the plasma parameters, material properties and device performance?
  - c) How Optical Emission Spectroscopy can be utilized for the end point detection of a plasma process? 3+4+3

[ 4 ]

4. The celebrated KdV equation describing weakly nonlinear long wave propagation in dispersive media reads as

$$\frac{\partial u}{\partial t} + 6u \frac{\partial u}{\partial x} + \frac{\partial^3 u}{\partial x^3} = 0, \text{ where } u \equiv u(x, t).$$

- Identify the roles the terms  $6u \partial u / \partial x$  and  $\partial^3 u / \partial x^3$  plays here.
- Show that the above KdV equation offers a single 'soliton' solution in the form:

$$u(x - Vt) = \frac{V}{2} \operatorname{sech}^2 \left[ \frac{\sqrt{V}}{2} (x - Vt) \right]$$

where  $V$  denotes the phase speed of the soliton.

- Hence point out the salient features of a soliton.

2+6+2

5. a) During the breakdown of gases, how the current is grown in the presence of secondary processes as per Townsend theory?
- b) What is Paschen's law? How the knowledge of Paschen's law can help you to get an idea about the breakdown condition in a glow discharge plasma within a vacuum chamber?
- c) Why plasma formation of gases is preferred for different material processing and other applications?

6+3+1

[ 3 ]

3. We focus on the macroscopic fluid description of plasmas. Together with Maxwell's equations, the fluid approach can almost provide a complete dynamic evolution of plasmas. Moreover, the fluid model can explain most of the laboratory experiments, thereby making it popular among the plasma physics community.

- Write down the basic 1D Maxwell-fluid equations that describe the dynamics of electron plasma oscillations in a cold collisionless unmagnetized plasma.
- Hence by employing the usual linearization procedure obtain the dispersion relation for the electron plasma oscillations.
- What do you mean by the 'cold plasma approximation' and the 'collisionless approximation' of plasmas.

3+5+2




Figure 1: Variation of the disposition rate with the Radio Frequency (RF) Power. [Refer to Question 6. (d)]

[ Turn over