

**B.E. ELECTRONICS AND TELE-COMMUNICATION FOURTH YEAR SECOND SEMESTER EXAM 2023**

**COLLOID AND SURFACE ENGINEERING**

**Time : 3 hours**

**Full Marks : 50**

**Part -I**

**Use Separate Answer scripts for each Part**

**Answer Any Two Questions**

**Clearly mention all the assumptions**

**Assume any missing data and mention it clearly**

1. a) Write short notes (any three)

[5x3=15]

- a) photo lithography
- b) Microcontact printing
- c) electron beam lithography
- d) Scanning probe induced oxidation

b) Explain in detail one gas-phase and one liquid-phase bottom-up process for 0-D nanomaterials fabrication.

[5x2=10]

2. a) Derive Poisson-Boltzman Equation. What is Debye-Huckel approximation.

[8+7]

b) A coating consists of consisting of 55 vol% TiO<sub>2</sub> (spherical, surface-treated pigment particles), 32 vol% emulsion, and 13 vol% water (containing some dissolved compounds). The oil content of the emulsion is 50 vol.%. The oil droplet and TiO<sub>2</sub> particle diameters are 1.2 and 0.3 μm, respectively. The temperature is 25°C and the viscosity of the medium (thickened aqueous phase) is 1 kg m<sup>-1</sup> s<sup>-1</sup>. The Hamaker constants of the oil and of TiO<sub>2</sub> in vacuum are  $5 \times 10^{-20}$  and  $15 \times 10^{-20}$  J, respectively. The Hamaker constant of the aqueous medium in vacuum is  $3.75 \times 10^{-20}$  J. The medium concentration of salt, NH<sub>4</sub>OH (applied to control pH to basic conditions), in the coating formulation is 10<sup>-5</sup> M. The relative permittivity of the medium is 80. The medium is water with a salt, NH<sub>4</sub>OH (1 : 1 salt) at 10<sup>-5</sup> M and  $\kappa^{-1}$  is calculated using the formula  $\kappa^{-1}=0.304/\sqrt{C}$ .

i) Calculate the attractive Van der Waals interaction energy between two oil droplets, between two TiO<sub>2</sub> particles, and between an oil droplet and a TiO<sub>2</sub> particle in the coating when the distance between the colloids (in all cases) is 15 nm. Comment on the result.

[ Turn over

ii) It has been decided to stabilize both pigments and oil droplets electrostatically. The surface potential for both colloids is  $-50$  mV. Calculate the electrostatic and composite (overall) interaction energies between an oil droplet and a  $\text{TiO}_2$  particle in the coating when the distance between them is  $10$  nm. Based on these calculations, what can you conclude more generally about the interaction of oil droplets and  $\text{TiO}_2$  particles in the coating? [10]

3. a) Derive an expression of heat of immersion relating surface tension. [7]

b) What is Deryaguin's approximation. How salt concentration affect interaction energies? [10+3]

c) The following data are available from a measurement of the zeta potential in an aqueous suspension of kaolin particles at  $25^\circ\text{C}$ :

- diameter of the spherical particles:  $0.75$  micrometre ( $\mu\text{m}$ );
- concentration of NaCl in water:  $0.003$  M;
- movement of the particle:  $350$   $\mu\text{m}$ ;
- time for movement of particle:  $4.2$  s;
- potential of the field in a  $10$  cm cell:  $200$  V.

Moreover, the following values are available for the  $f$  correction parameter of the Henry equation for various values of  $\kappa R$  ( $\kappa^{-1}$  is the Debye length and  $R$  is the radius of the particle):

$\kappa R$	$f(\kappa R)$	$\kappa R$	$f(\kappa R)$
0	1.000	5	1.160
1	1.027	10	1.239
2	1.066	25	1.370
3	1.101	100	1.460
4	1.133	$\infty$	1.500

i) Calculate the electrophoretic mobility of the colloid particles.

ii) Provide an estimation of the zeta potential of the particles. [5]

**B.E. ELECTRONICS AND TELE-COMMUNICATION ENGINEERING FOURTH YEAR  
SECOND SEMESTER EXAM 2023**

**COLLOID AND SURFACE ENGINEERING**

Part-II

FM-50

**Use Separate Answer script for each part**

**Assume any missing data**

1. (i) Calculate the electrophoretic mobility of a 50 nm diameter spherical colloid particle in an aqueous solution of NaCl at 298 K. The  $\xi$ -potential is 0.02 V. The concentration of NaCl in the solution is 100 mol/m<sup>3</sup>. Given: The Debye length is ~1 nm at this concentration of the salt. Given:  $\epsilon = 80$ . (7)
- (ii) What is streaming potential? How is it developed? (1+2)
- (iii) Write short notes on (any two): Peclet Number, Krafft Point, Donnan Equilibrium (10)
  
2. (i) The aggregation number of the surfactant C<sub>10</sub>H<sub>21</sub>N(CH<sub>3</sub>)<sub>3</sub>Br has been reported to be 36. Can its micelle be spherical? (7)

or

The variation of osmotic pressure of a polystyrene solution in toluene with its concentration at 298 K is given below.

$c$ (kg/m <sup>3</sup> )	0.2	0.4	0.6	0.8	0.9
$\pi_o$ (cm of toluene)	0.04	0.09	0.16	0.22	0.28

Determine the molecular weight of the polymer from these data. Given: density of toluene at 298 K = 860 kg/m<sup>3</sup>.

- (iii) How the critical micelle concentration depends upon electrolyte concentration and pH of the solution? (4)
- (iv) State Fowkes modification over Girifalco and Good Correlation regarding interfacial tension. (4)
  
3. (i) Discuss how the pressure difference to the shape of the planar, cylindrical and spherical surface. (10)
- (ii) Write down the Assumptions made in BET adsorption isotherm. Differentiate chemisorption from physisorption. (3+2)